

Banjo Forte 400 SC

The best weapon against black pod disease

ADAMA

New innovative fungicide for the control of Black pod disease

PERFECT COVERAGE - FAST ACTING AND LONG LASTING

Banjo Forte 400 SC is a non-copper foliar fungicide with protective and translaminar mode of action.

Banjo Forte 400 SC Fluazinam Dimethomorph ADAMA FUNGICIDE **BANJO FORTE** Preventative and Systemic activity 400 SC anti sporulating 200g/I dimethomorph + 200g/l fluazinam activity ADAMA uspension concentrate Effective non-copper fungicide the Control Black pod Disease on Coc No. Date of Manuf: Expiry date: 250m Makhteshim Ltd, Adama West Africa Ltd Zone, P.O. Box Sheva 84100, Isra CAUTION



BENEFITS FOR THE FARMER



Easy to use- SC formulation





Resistance Effective Management-The benefit of two active Ingredients



Can be tankmixed with foliar fertilizers and insecticides



Non copper no heavy metal added



Excellent rain fastness-Only 30 minutes

Banjo Forte 400 SC

is used as a protectant and curative fungicide to control black pod disease on cocoa.

DIRECTION FOR USE:

Mixing and application half fill the spray tank with clean water. Add the recommended amount of Banjo Forte 400 SC and water while agitating the spraying machine continuously. Spray to drip on pods.

Always read the label carefully before use





Find us on Facebook ADAMA West Africa

Crop	Disease	Application rate
Сосоа	Black pod	1,125 L in 225L of water / ha
	(le Phytophtora megakaria and phytophtora palmivora)	450 L in 90L of water/ ha
		75 ml in 15L of water/ Knapsack

WHO Class III

Timing and frequency of application: Apply at 4 weekly intervals from April/May (depending on the time of the rains) to September/November annually.

Pre-harvest interval: 7 days **Re-entry period:** at least 24 hrs after treatment.

Manufactured by:



Adama West Africa Ltd. No. 4,4th Norla Street, North Labone Accra, Ghana Tel.: 0544325030 Info.westafrica@adama.com www.adama.com/west-africa

ADAMA