


**Mastercop®** is a new fungicide and bactericide of the copper group, developed with a **unique formulation** of outstanding quality, that delivers an excellent control of Coffee Berry disease, Coffee Leaf Rust and Bacterial Blight of Coffee.



## Mastercop is a game changer of the coppers

Powder Coppers	Mastercop®
Large particles size thus reduced absorption by the cuticle	Very small particle size thus higher efficacy compared to all other
Blue or coloured residue on the Crop. Low market quality of the fruits and reduced photosynthesis activity	No discolouration of the crop. High market quality of crop and more photosynthetic activity
Easily washed by irrigation or rain	Faster absorption thus reduced wash off
Nozzle wear	No sedimentation and nozzle wear
High rate of use thus risk of phytotoxicity and increase soil loading of copper	Low rate of use thus prevents phytotoxicity



**Mastercop®** is a preventive fungicide and bactericide containing copper sulphate pentahydrate, developed with a unique formulation of outstanding quality for the control of Coffee Leaf Rust and Berry Disease, early and late blight in tomatoes; halo blight, angular leaf spot, anthracnose and rust in French beans; black rot, black leg and soft rot in Broccoli.

**Direction for use:** Mix with water at the recommended rates and foliar apply. Fill the mix tank with water to half volume. Add the recommended quantity of Mastercop® 60SC. Fill the mix tank to full volume and mix well before application.

Crop	Disease	Rate/ha in 1000 L of water	Rate in 20 L of water	Repeat applications
Coffee	Coffee Berry	2 L	40 mL	Spray once in 4 weeks
	Coffee Leaf Rust	2 L	40 mL	Spray once in 3 weeks
Tomato	Early and Late Blight	1 L	20 mL	4 applications every 10 days
French beans	Halo Blight, Angular leaf spot, Anthracnose and Rust	1 L	20 mL	3 applications every 10 days
Broccoli	Black Rot, Black Leg and Soft Rot	1 L	20 mL	3 applications every 10 days

Pre-harvest interval for tomatoes, French beans and broccoli: 0 days

**Always read the label carefully before use**

**Global uses of Mastercop®**

Crop	Disease	Rate (L/Ha)
Corn	Bacterial Stalk Rot	0.6-1.5
Potato	Early and Late Blight	1.0-1.5
Wheat, Barley, Oats	Helminthosporium Spot, Blotch, Septoria Leaf Blotch	0.5-1.0
Berries	Bacterial Canker	1.5-3.0
	Fruit Rot, Phomopsis Twig Blight	1.0-2.0
Avocado	Anthracnose, Blotch, Scab	2.0-4.0
	Coffee Berry Disease (Colletotrichum coffeanum)	2.0-2.5
Coffee	Bacterial Blight (Pseudomonas syringae)	2.0-3.0
	Leaf Rust (Hemileia vastatrix)	1.0-1.5
Mango	Anthracnose	1.5-2.0
Bean	Brown Spot, Common Blight, Halo Blight	0.5-1.0
Carrot	Alternaria Leaf Spot, Cercospora Leaf Spot	0.5-1.5
Brassica	Black Leaf Spot, Black Rot, Downy Mildew	0.5-1.0
Cucurbits	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Gummy Stem Blight,	0.5-1.0
Eggplant	Powdery Mildew, Water-melon Bacterial Fruit Blotch (suppression)	0.5-1.5
	Alternaria Blight, Anthracnose, Phomopsis	
Okra	Anthracnose, Bacterial Leaf Spot, Leaf Spots, Pod Spot, Powdery Mildew	0.5-1.5
Onion, Garlic	Bacterial Blight, Downy Mildew, Purple Blotch	0.5-1.5
Pea	Powdery Mildew	0.5-1.0
Pepper	Anthracnose, Bacterial Leaf Spot, Cercospora Leaf Spot	0.5-3.0
Tomato	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight Gray Leaf Mold, Late Blight, Septoria Leaf Spot	1.0-3.0
Grapes	Black Rot, Downy Mildew, Phomopsis	1.5-3.0
	Canker Suppression	1.0-3.0
Citrus	Algal Spot, Melanose, Scab	1.5-5.0
	Alternaria Brown Rot	1.5-3.5
	Greasy Spot, Pink Pitting	0.5-2.5
	Phytophthora Brown Rot, Septoria Spot	1.5-3.5
Herbs	Downy Mildew	0.5-1.0
	Anthracnose	1.0-3.0
Macadamia	Phytophthora Blight (P. capsici), Raceme Blight (Botrytis cinerea)	1.0-2.0
Papaya	Anthracnose	1.5-3.0
Passion Fruit	Anthracnose	1.5-3.0
Roses	Downy Mildew, Powdery mildew, Botrytis	1.5-2.0
	Crown gall	2.0-2.5
Carnation	Carnation rust and Ringspot	1.0-1.5
	Fusarium and other soil-borne diseases	2.0-3.0

**Mastercop®** is omri listed. it may be used in certified organic production or food processing and handling according to the usda national organic program rule.

Manufactured by:



ADAMA

For further information please contact:

Robert Wachira

Tel: +254 713 463 937

www.adama.com/east-africa

Distributed by:



CropTech (K) LTD

P.O. Box 137-00618

Ruaraka, Nairobi

Tel. +254 722 703 911

www.croptechltd.com

**ADAMA**



Find us on Facebook  
ADAMA East Africa



# Mastercop®

More than just another copper



FUNGICIDE /BACTERICIDE

- Very good bactericidal and fungicidal effect
- Rain-fastness
- Translaminar activity
- Absence of color, therefore photosynthetic efficiency
- Low application rates per Ha
- Low copper loading to the soil
- Liquid formulation easy to handle
- Does not wear or block the spray equipment

Simply. Grow. Together.

ADAMA.COM